

DECREASED EVENT-RELATED THETA POWER AND PHASE-SYNCHRONY IN
YOUNG BINGE DRINKERS DURING TARGET DETECTION: AN
ANATOMICALLY-CONSTRAINED MEG APPROACH.

Authors: Ángeles Correas, Eduardo López-Caneda, Lauren Beaton, Socorro Rodríguez Holguín, Luis Miguel García-Moreno, Luis Fernando Antón Toro, Fernando Cadaveira, Fernando Maestú, Ksenija Marinkovic

This is the peer reviewed version of the following article: Correas A; López-Caneda E; Beaton L; Rodríguez Holguín S; García-Moreno LM; Antón Toro LF; Cadaveira F; Maestú F; Marinkovic K (2018). Decreased event-related theta power and phase-synchrony in young binge drinkers during target detection: An anatomically-constrained MEG approach. *Journal of Psychopharmacology*, First Published online October 24, 2018.
doi: [10.1177/0269881118805498](https://doi.org/10.1177/0269881118805498).

This article may be used for non-commercial purposes in accordance with SAGE publications Terms and Conditions for Use of Self-Archived Versions

**Decreased event-related theta power and phase-synchrony
in young binge drinkers during target detection: An
anatomically-constrained MEG approach.**

Journal:	<i>Journal of Psychopharmacology</i>
Manuscript ID	JOP-2018-3460.R1
Manuscript Type:	Original Paper
Date Submitted by the Author:	06-Aug-2018
Complete List of Authors:	Correas, Angeles; San Diego State University, Psychology López-Caneda, Eduardo; University of Minho, Neuropsychophysiology Lab, Research Center on Psychology Beaton, Lauren; San Diego State University, Psychology Rodríguez Holguín, Socorro; Universidade de Santiago de Compostela Facultade de Psicoloxia, Clinical Psychology and Psychobiology García-Moreno, Luis Miguel; Complutense University of Madrid, Department of Psychobiology and Methodology in Behavioral Sciences Antón Toro, Luis Fernando; Centre of Biomedical Technology, Laboratory of Cognitive and Computational Neuroscience (UCM-UPM) Cadaveira, Fernando; Universidade de Santiago de Compostela Facultade de Psicoloxia, Clinical Psychology and Psychobiology Maestú, Fernando; Centre of Biomedical Technology, Laboratory of Cognitive and Computational Neuroscience (UCM-UPM); Complutense University of Madrid, Department of Experimental Psychology; Network of Center for Biomedical Research, CIBER-BBN Marinkovic, Ksenija; San Diego State University, Psychology; University of California San Diego, Radiology
Please list at least 3 keywords which relate to your manuscript::	Binge drinking, alcohol, magnetoencephalography, theta oscillations, attention
Abstract:	<p>Background: The prevalence of binge drinking (BD) has been on the rise in recent years. It is associated with a range of neurocognitive deficits among adolescents and young emerging adults who are especially vulnerable to alcohol use. Attention is an essential dimension of executive functioning and attentional disturbances may be associated with hazardous drinking. The aim of the study was to examine the oscillatory neural dynamics of attentional control during visual target detection in emerging young adults as a function of BD.</p> <p>Method: Fifty-one first-year university students (18±0.6 years) were assigned to light drinking (LD, N=26), and BD (N=25) groups based on their alcohol consumption patterns. High-density magnetoencephalography (MEG) signal was combined with structural magnetic resonance imaging (MRI) in an anatomically-constrained MEG model to estimate event-related source power in theta (4-7 Hz) frequency band. Phase-locked co-</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>oscillations were further estimated between the principally activated regions during task performance.</p> <p>Results: Overall, the greatest event-related theta power was elicited by targets in the right inferior frontal cortex (rIFC) and it correlated with performance accuracy and selective attention scores. BDs exhibited lower theta power and dysregulated oscillatory synchrony to targets in the rIFC which correlated with higher levels of alcohol consumption.</p> <p>Conclusions: These results confirm that a highly interactive network in the rIFC subserves attentional control, revealing the importance of theta oscillations and neural synchrony for attentional capture and contextual maintenance. Attenuation of theta power and synchronous interactions in BDs may indicate early stages of suboptimal integrative processing in young, highly functioning BDs.</p>
--	--



TITLE:

Decreased event-related theta power and phase-synchrony in young binge drinkers during target detection: An anatomically-constrained MEG approach.

AUTHOR NAMES AND AFFILIATIONS:

Angeles Correas^a, Eduardo López-Caneda^b, Lauren Beaton^a, Socorro Rodríguez Holguín^c, Luis Miguel García-Moreno^d, Luis F. Antón-Toro^e, Fernando Cadaveira^c, Fernando Maestú^{e,f,g}, Ksenija Marinkovic^{a,h}.

^a Department of Psychology, San Diego State University, San Diego, USA.

^b Neuropsychophysiology Lab, Research Center on Psychology (CIPsi), School of Psychology, University of Minho, Braga, Portugal.

^c Department of Clinical Psychology and Psychobiology, University of Santiago de Compostela, Spain.

^d Department of Psychobiology and Methodology in Behavioral Sciences, Complutense University of Madrid, Madrid, Spain.

^e Laboratory of Cognitive and Computational Neuroscience (UCM-UPM), Centre of Biomedical Technology (CTB), Madrid, Spain.

^f Department of Experimental Psychology, Complutense University of Madrid, Spain.

^g Network of Center for Biomedical Research (CIBER-bbn), Madrid, Spain.

^h Department of Radiology, University of California at San Diego, San Diego, USA.

Please address correspondence to:

Ksenija Marinkovic Ph.D., Department of Psychology, 5500 Campanile, Dr., San Diego State University, San Diego, CA 92182, USA.

Email: kmarinkovic@mail.sdsu.edu

Telephone number: 619-594-1510

Abstract: 249 words

Manuscript: 4979 words

References: 163

Figures: 4

Tables: 2

ABSTRACT

Background: The prevalence of binge drinking (BD) has been on the rise in recent years. It is associated with a range of neurocognitive deficits among adolescents and young emerging adults who are especially vulnerable to alcohol use. Attention is an essential dimension of executive functioning and attentional disturbances may be associated with hazardous drinking. The aim of the study was to examine the oscillatory neural dynamics of attentional control during visual target detection in emerging young adults as a function of BD.

Method: Fifty-one first-year university students (18 ± 0.6 years) were assigned to light drinking (LD, $N=26$), and BD ($N=25$) groups based on their alcohol consumption patterns. High-density magnetoencephalography (MEG) signal was combined with structural magnetic resonance imaging (MRI) in an anatomically-constrained MEG model to estimate event-related source power in theta (4-7 Hz) frequency band. Phase-locked co-oscillations were further estimated between the principally activated regions during task performance.

Results: Overall, the greatest event-related theta power was elicited by targets in the right inferior frontal cortex (rIFC) and it correlated with performance accuracy and selective attention scores. BDs exhibited lower theta power and dysregulated oscillatory synchrony to targets in the rIFC which correlated with higher levels of alcohol consumption.

Conclusions: These results confirm that a highly interactive network in the rIFC subserves attentional control, revealing the importance of theta oscillations and neural synchrony for attentional capture and contextual maintenance. Attenuation of theta power and synchronous interactions in BDs may indicate early stages of suboptimal integrative processing in young, highly functioning BDs.

KEYWORDS: Binge drinking, alcohol, magnetoencephalography, theta oscillations, attention

1. INTRODUCTION

Binge drinking (BD) is highly prevalent among adolescents and young adults in most Western countries (Eurobarometer, 2010; Johnston, 2011; SAMHSA, 2013) and is associated with a wide range of negative social and health consequences (Hingson et al., 2017; Hingson and White, 2013). This pattern of consumption is characterized by intake of intoxicating quantities of alcohol in a short time interval interspersed with periods of abstinence (Courtney and Polich, 2009). BD is usually defined as imbibing five/four or more drinks for men/women within a two-hour interval, which is likely to bring blood alcohol concentration (BAC) to 0.08% or above (National Institute of Alcohol and Alcoholism, 2004). However, these levels are commonly exceeded by emerging young adults who engage in bouts of much higher levels of alcohol intake (Naimi et al., 2010; Patrick and Terry-McElrath, 2017; Terry-McElrath and Patrick, 2016).

Adolescence emerging into adulthood is a protracted but critical period of neurodevelopment characterized by profound changes (Sherman et al., 2014; Simmonds et al., 2014; Walhovd et al., 2012) which mainly involve prefrontal cortex (PFC) and other higher-order association areas (Gogtay et al., 2004; Lebel and Beaulieu, 2011) and contribute to refinement of cognitive efficiency (Casey et al., 2005). As it continues to mature well into mid 20's, the adolescent brain is more sensitive to the impact of ethanol exposure, with the frontal lobes being particularly vulnerable (Cservenka and Brumback, 2017; Hermens et al., 2013; Jacobus and Tapert, 2013; Silveri et al., 2016; Squeglia et al., 2014). Extensive evidence obtained from rodent models indicates that BD during adolescence is a high risk factor for the development of alcohol-use disorders (AUD) and enduring neural changes (Crabbe et al., 2011; Crews et al., 2016).

Several studies have reported that BDs differ from light drinkers (LDs) on EEG measures in the absence of behavioral differences (Folgueira-Ares et al., 2017; López-Caneda et al., 2012, 2013). However, other studies report that BDs tend to have lower scores than light-drinkers (LDs) on tasks probing high-level cognitive abilities such as memory and executive functions (for reviews, see Stephens and Duka, 2008; Jacobus and Tapert, 2013; Petit et al., 2014). Attention is a key component of executive functions and is involved in minimizing distraction and in optimal allocation of cognitive resources to task-relevant stimuli (Diamond, 2013). Impairments of attention are related to increased impulsivity and may contribute to engaging in

1
2
3 hazardous drinking (Crews and Boettiger, 2009; de Wit, 2009). Despite its importance for
4 response inhibition and cognitive control, attention has been scarcely explored in BD
5 population. Lannoy et al. (2017) used the Attention Network Task (Fan et al., 2005) and
6 reported impairments of alerting and executive control dimensions of attention in BDs. Using the
7 same task, Maurage et al. (2014) showed that alcohol-dependent individuals presented
8 attentional deficits specifically in the executive control of attention compared with social
9 drinkers. These results suggest that attentional disturbances might have a key role in the
10 development of AUD in the context of broader cognitive impairments (Muller-Oehring and
11 Schulte, 2014; Oscar-Berman and Marinković, 2007; Sullivan and Pfefferbaum, 2005).
12 Convergence of neuroimaging and human lesions studies has indicated that the right-dominant
13 fronto-parietal network is crucially involved in attentional processes (Klinke et al., 2017; Vossel
14 et al., 2014). Functional MRI (fMRI) studies have confirmed engagement of a right-lateralized
15 network during target detection with a nexus in the right inferior prefrontal cortex (Gitelman et
16 al., 1999; Nobre et al., 1997; Shulman et al., 2010; Vossel et al., 2014).

17
18
19 A previous EEG study has reported alterations in event-related oscillations (ERO) in BDs during
20 the same equiprobable Go/NoGo task as the one employed in the present study (López-
21 Caneda et al., 2017). This study found lower delta and theta power during target detection in
22 BDs. They proposed that this power decrease in low frequencies may reflect impairments in the
23 neural circuit involved in both the activation and inhibition of a response. Moreover, ERO studies
24 in individuals with AUD have reported attenuated power in slow frequencies (delta and/or theta)
25 in alcoholics during Go/NoGo task (Colrain et al., 2011) and equiprobable Go/NoGo task
26 (Kamarajan et al., 2004; Pandey et al., 2016; Porjesz et al., 2005) which has been interpreted
27 as a deficiency in inhibitory and attentional processing.

28
29
30 Event-related theta (ERT) is sensitive to cognitive effort as it increases in response to higher
31 demands of cognitive control (Brier et al., 2010; Cavanagh and Frank, 2014; Hanslmayr et al.,
32 2008; Jensen and Tesche, 2002; Rosen et al., 2016). Furthermore, ERT is strongly modulated
33 by alcohol. Acute alcohol intoxication attenuates ERT especially on trials evoking cognitive
34 interference (Beaton et al., 2017; Kovacevic et al., 2012; Marinkovic et al., 2012; Rosen et al.,
35 2016), indicating that alcohol primarily affects executive functions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Theta oscillations have also been implicated in integrating task-relevant representations with top-down effects across long-range neural envelopes (Halgren et al., 2015; Hasselmo and Stern, 2014; Wang et al., 2005). Phase-locking values (PLV) are a measure of phase consistency between two signals (Lachaux et al., 1999) and are suitable for investigating co-oscillations between different brain areas at a level of an interactive dynamic network (Varela et al., 2001). It has been shown that PLVs between the lateral and medial prefrontal cortical areas are sensitive to cognitive conflict (Beaton et al., 2017) and that acute intoxication dysregulates those interactions in humans and rodents (Beaton et al., 2017; Ehlers et al., 2012). There is a paucity of studies assessing task-induced functional connectivity in BDs with temporally sensitive methods. However, altered functional connectivity during resting-state has been reported in BDs compared to LDs (Correas et al., 2015; Correas et al., 2016).

The aim of the present study was to examine neural dynamics of the attentional circuitry during target detection in adolescents emerging into adulthood as a function of BD. To this purpose we employed an anatomically constrained MEG method (aMEG) which is a multimodal approach that combines whole-head MEG recordings and high-resolution MRI images within a distributed source model (Dale et al., 2000; Lin et al., 2004; Marinkovic et al., 2003). It allows for the estimation of *where* the oscillatory changes are occurring, and it provides a highly precise insight into the temporal sequence ("*when*") of the involved neural components. Maps of event-related power estimates and the associated time courses of source estimates were analyzed across time and cortical space in time-frequency domain (Kovacevic et al., 2012; Lin et al., 2004; Marinkovic et al., 2012; Rosen et al., 2016). Additionally, co-oscillations between the principally activated cortical regions provided insight into attentional function in real time and at the level of an interactive system in BD and LD groups (Beaton et al., 2017).

2. MATERIALS AND METHODS

2.1 Participants

Fifty-one first-year students of the Complutense University of Madrid (Madrid, Spain) participated in the study. They were assigned to a binge drinking (BD) (N=25, 13 females) or to a light drinking (LD) group (N=26, 14 females) based on a questionnaire and a semi-structured interview inquiring about their alcohol and other drug use. Participants provided a record of their daily alcohol consumption indicating the type(s) and the quantity of the beverage(s) they consumed in the past month as well as the length of time (in hours) it took them to imbibe these beverages (Table 1). Their estimated Blood Alcohol Concentration (*est*-BAC) was calculated for each drinking episode based on the information they provided for the past month, as well as their gender and weight, according to the following algorithm (Windmark, 1981):

$$\text{estimated BAC} = \left(\frac{G \text{ alcohol}}{W \times bw} \right) - mr \times \text{hours}$$

where G corresponds to the highest level of alcohol (grams) consumed on one occasion in the previous month; W is the body weight (Kg); bw or body water is a constant related to the water content of the human body with a value of 0.68/0.55 for male/female; *mr* is the metabolism rate with a value of 0.15/0.18 for male/female. We considered the *est*-BAC as a rough index of alcohol intoxication for each subject. Participants reaching an *est*-BAC of 0.08% or above at least once during the previous month were classified as BDs. In contrast, the control group consisted of students who did not achieve that BAC in the past month. It is important to note that the legal drinking age in Spain is 18 years of age.

Participants completed personality tests related to impulsivity variables including Barrat Impulsivity Scale (BIS) (Patton et al., 1995) and Sensation Seeking Scale (SSS) (Zuckerman and Link, 1968). Their selective and sustained attention was measured with the D2 attentional test (Brickenkamp and Seisdodos, 2009), a neuropsychological test in which participants are asked to cross out any "d" letter. Also, their sensitivity to reward was assessed by means of the Iowa Gambling Task (IGT) (Bechara et al., 1994).

Prospective participants with family history of alcohol abuse were excluded from the study as well as individuals reporting personal history of psychiatric disorders based on DSM-IV-TR

criteria (American Psychiatric Association, 2000) and/or psychopathological traits as assessed by the Symptom Checklist-90 Revised questionnaire (Derogatis and Melisaratos, 1983). Subjects reported no current medical condition and were medication free at the time of the study. They also completed the Alcohol Use Disorders Identification Test (AUDIT) (Guillamón et al., 1999) and subjects who scored 20 or above were excluded for having AUD. Tobacco smoking was a controlled variable. According to the Syndrome Nicotine Dependence Scale (Becoña et al., 2011), four participants fulfilled the criterion of nicotine dependence, three BDs and one LD.

Participants were asked to refrain from alcohol consumption for at least 24 hours prior to data acquisition. All of them provided written informed consent for the study and were monetarily reimbursed for their participation. The study was approved by the Ethics Committee of the Complutense University of Madrid. Principles of Declaration of Helsinki were followed.

Table 1. Demographic, drinking, personality and neuropsychological variables.

	<i>LDs</i>	<i>BDs</i>	<i>U values</i>
<i>N (females)</i>	26 (14)	25 (11)	164(χ^2)
<i>Age</i>	18±0.8	18±0.4	650
<i>Est BAC</i>	0.017±0.02	0.17±0.072	702**
<i>Length of drinking (mos)</i>	4.55±3.71	20.67±10.45	205.5**
<i>Age of drinking onset</i>	16.6±1.01	14.92±1.08	83*
<i>Drinking days past mo.</i>	1.5±1.8	7.82±2.2	184.5**
<i>Drinks on a regular day</i>	1.1±0.5	4±1.7	267.5**
<i>Drinks on a heavy day</i>	2.4±1.1	5.8±2.6	248.5**
<i>Drunk days past mo.</i>	0	4.84±2.56	688.5**
<i>Total AUDIT score</i>	1.7±1.3	7.7±1.9	648**
<i>Sens. Seeking –Tot.</i>	16.29±3.5	21.48±3.53	257**
<i>Thrill & Adventure</i>	5.76±2.66	7.58±2.35	278
<i>Disinhibition</i>	2.48±1.22	5.17±1.58	322**
<i>Experience Seeking</i>	5.56±1.58	5.43±1.77	175
<i>Boredom Suscept</i>	2.54±1.81	3.17±1.19	222
<i>Barrat Impuls. - Tot.</i>	38.87±10.6	44.38±11.5	241
<i>Emotional</i>	14.42±3.81	12.84±3.95	261.5
<i>Motor</i>	14.77±6.55	12.96±5.53	320
<i>Non-planning</i>	15±4.53	14.2±4.95	242
<i>Attention D2 - Total</i>	498.23±64.1	468.84±75.26	157
<i>IGT – total earnings</i>	12.89±21.85	11.15±20.5	216
<i>IGT – total losses</i>	19.03±17.25	21.6±29.3	204
<i>Sleeping hours previous night</i>	7.2±0.98	7.04±1.17	257.5

Mean±SD are listed for all variables. Group comparisons were calculated with a nonparametric Mann-Whitney U test except for age which was tested with χ^2 . LDs: light drinkers; BDs: binge drinkers; estBAC: the highest estimated Blood Alcohol Concentration reached in the past month; AUDIT: Alcohol Use Disorders Identification Test; IGT: Iowa Gambling Task. Significance level is indicated as follows: *p < 0.05, **p < 0.01, ***p < 0.001

2.2 Task

Participants performed a target detection task with equiprobable target (Go) and nontarget (NoGo) stimuli which consisted of blue/green squares/circles. Participants were instructed to fixate on a cross in the center of the screen and to press a button as fast as possible to targets (green circles and blue squares) and to withhold responding to non-target (blue circles and green squares). The stimuli were presented for 100 ms with a stimulus onset asynchrony (SOA) of $1,400 \pm 200$ ms within a visual angle of 3.4 degrees. They were presented in random order in two blocks of 200-225 stimuli (Figure 1). Half of subjects pressed the button with the left and half with the right hand in a counterbalanced manner.

[insert Figure 1.]

2.3 Data acquisition and analysis

2.3.1 MRI

Structural MRI images were used to define the model for the volume conductor and the solution space for MEG source localization analysis. Images were acquired with a General Electric 1.5 Tesla using an eight-channel head coil. The imaging protocol consisted of: 3D T1-weighted high-resolution images using a Fast-Spoiled Gradient Echo sequence [TR/TE/TI=11.2/4.2/450 ms; flip-angle= 12°; FoV= 250 mm; acquisition matrix= 256 x 256; slice thickness=1 mm]. Structural images were used to reconstruct each person's cortical surface with FreeSurfer software. Inner skull surface was used for a boundary element model of the volume conductor in the forward calculations. The solution space was approximated by ~5000 free-rotation dipoles along the gray-white matter surface in the cortex, with spacing between dipole locations ~7 mm (Kovacevic et al., 2012).

2.3.2 MEG

High-density MEG signals were recorded from 204 channels (102 pairs of planar gradiometers) with a whole-head Neuromag Vectorview system (Elekta) in a magnetically and electrically shielded room. The signals were recorded continuously with a 1000 Hz sampling rate and

1
2
3 filtered online with a band pass filter 0.1-330 Hz. The position of magnetic coils attached to the
4 skull, the main fiducial points, as well as a large array of random points spread across the
5 scalp were digitized with 3Space Isotrak II system for subsequent precise co-registration with
6 structural MRI images. Trials with incorrect responses were excluded from the analysis. The
7 number of included trials was equated across the target and nontarget task conditions for each
8 subject. The data were analyzed with a time-frequency analysis stream that uses MATLAB
9 routines (Beaton et al., 2017; Kovacevic et al., 2012; Marinkovic et al., 2012; Rosen et al., 2016)
10 that incorporates publicly available packages including Fieldtrip (Oostenveld et al., 2011),
11 EEGLab (Delorme and Makeig, 2004) and MNE (Gramfort et al., 2014). Continuous data were
12 bandpass filtered from 0.1 to 100 Hz and epoched from -600 to 1000 ms relative to stimulus
13 onset for the stimulus-locked analysis. For each epoch, the data were downsampled by a factor
14 of 4 to 250 Hz. Epoched data were then passed through automatic threshold rejection to
15 remove trials that were contaminated with artifacts. Independent component analysis was used
16 to remove eye-blinks and heart beat artifacts (Delorme and Makeig, 2004). Complex power
17 spectrum was calculated for each trial using convolution with complex Morlet wavelets (Lachaux
18 et al., 1999) in 1 Hz increments from 4 to 7 Hz for theta band. The first and last 300 ms of each
19 epoch were discarded to remove edge artifacts resulting from wavelet analysis. Wavelet results
20 were visually inspected across all epochs for any additional artifacts. We used an anatomically
21 constrained MEG (aMEG) source modeling which comprises cortically constrained minimum
22 norm estimates of the event-related theta power (Dale et al., 2000; Lin et al., 2004). To estimate
23 the noise covariance for calculation of the inverse and to prevent biasing the inverse solution,
24 empty room data were detrended and band-pass filtered between 3 and 40 Hz. The signal-to-
25 noise ratio equaling 5 (Lin et al., 2004; Marinkovic, 2003) was used for scaling of the noise
26 covariance matrix in calculation of the inverse operator. The identity matrix was used for noise-
27 sensitivity normalization of the source-space solution. The noise-sensitivity normalized
28 estimates of total source power were obtained at each location on the cortical surface at each
29 frequency. Source power estimates were calculated based on cortically constrained minimum
30 norm estimates (Kovacevic et al., 2012; Lin et al., 2004; Marinkovic et al., 2012). For each
31 subject, a map of total source power for theta was estimated by averaging across theta
32 frequency (4-7 Hz) and across all trials for each condition. ERT power was baseline-corrected
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 by subtracting the mean theta source power estimate in the 300 ms prestimulus period and
4 expressed as percent signal change from baseline. Uncorrected theta power in the baseline did
5 not differ between groups ($F(1,50) = 0.02, p = 0.86$) or conditions ($F(1,50) = 0.7, p = 0.37$)
6 confirming that the stimulus-evoked changes were not affected by possible differences in the
7 baseline. Intersubject averages were created by morphing each subject's reconstructed surface
8 onto an average representation after aligning their cortical sulcal-gyral patterns (Fischl et al.,
9 1999) and averaging individual source power estimates.

10
11
12
13
14
15
16 Region-of-interest (ROI) analysis was conducted to further examine possible interactions of task
17 condition, group, and gender on event-related changes in theta power. Unbiased ROIs were
18 selected based on the overall group average across all subjects for each task condition and
19 comprised dipole locations along cortical surface with most notable source power. The same set
20 of group-based ROIs was used for all subjects in a manner blind to their individual activations by
21 applying an automatic spherical morphing procedure (Fischl et al., 1999). Early visual activity
22 was assessed in the occipital region (Occ) while the anterior (aIFC) and posterior inferior frontal
23 cortices (pIFC) and sensorimotor cortices (sMOT) were examined bilaterally as shown in Figure
24 1. Additionally, co-oscillatory interactions between cortical ROIs were estimated by calculating
25 phase-locking values (PLV) between pairs of frontal ROIs in each hemisphere (Figure 3)
26 (Beaton et al., 2017; Lachaux et al., 1999).

27
28
29
30
31
32
33
34
35
36 A nonparametric cluster-based permutation test (CBPT)(Maris and Oostenveld, 2007), was
37 used to test for significant group differences in event-related theta power, as well as in PLVs. At
38 each time step, group differences were tested by means of a 1-way ANOVA test which provided
39 F values used in the CBPT. Time clusters consisted of ≥ 25 contiguous time points yielding
40 100ms time windows which systematically showed group difference with a significance
41 threshold $p \leq 0.05$. Cluster-statistics were assessed through the sum of all F-values
42 corresponding to the time steps comprised in cluster. This procedure was repeated 5000 times
43 after shuffling the original group's labels which created a null distribution for each comparison.
44 The cluster-statistics obtained from the original data set were compared with those in the
45 randomized data, while keeping the maximum statistic at each repetition. The resulting CBPT p-
46 value represented the proportion of the permutation distribution with cluster-statistic value of the
47
48
49
50
51
52
53
54
55
56
57
58
59
60

original data. Only the time-cluster that survived the CBPT at $p < 0.05$ was used to define time windows for the subsequent analyses.

No significant effects of Gender were observed in any of the analyses. Nonparametric Spearman's rho coefficient was used to compute correlations.

3. RESULTS

3.1 Behavioral Performance

Behavioral performance was assessed by measuring accuracy in detecting targets and non-targets and reactions times to targets. Behavioral results are summarized in Table 2. BDs showed a marginal tendency to respond less accurately and with longer RTs in comparison to LDs.

Table 2. Behavioral data for the LD and BD groups (mean \pm SD).

Behavioral Performance	LDs	BDs	t-test (49)	P-values
Reaction Time (ms)	509.08 \pm 73.61	531.46 \pm 74.32	1.03	$p = 0.1$
% correct Responses (target)	94.59 \pm 4.5	92.01 \pm 5.41	1.84	$p = 0.09$
% correct Non-Responses (non-target)	88.66 \pm 7.8	86.76 \pm 9.14	0.79	$p = 0.45$

3.2 MEG: Event-related theta

The aMEG method provided insight into spatio-temporal stages of the ERT power during target detection. As shown in Figure 1, there is an early theta increase estimated to the occipital cortex across both groups and conditions. However, subsequent theta increase was observed only to the target stimuli. In contrast, nontargets did not elicit appreciable ERT increase relative to the baseline. The greatest theta activity overall was observed in the right IFC peaking after ~450 ms. Both the right aIFC and pIFC were also sensitive to group differences with BDs showing attenuated theta to targets as compared to LDs (Table 3) during the 380-510 ms time window.

[insert Figure 2.]

Table 3. MEG Theta Band Results.

ROIs	Condition, F(1,50)	Group, F(1,50)	Condition x Group, F(1,50)	Target BD vs LD, F(1,50)	NonTarget BD vs LD, F(1,50)
<i>Right aIFC</i>	41.65***	4.1*	4.74*	4.74*	0.006

<i>Right pIFC</i>	41.06***	4.12*	4.18*	4.34*	0.004
<i>Right sMot</i>	21.25***	3.8	0.7	4.02	2.3
<i>Left aIFC</i>	30.1***	0.5	1.82	1.33	0.5
<i>Left pIFC</i>	28.68***	1.94	0.6	0.75	0.15
<i>Left sMot</i>	20.54***	0.7	0.003	0.14	1.27
<i>Occipital</i>	0.97	0.31	0.001	0.02	0.03

ANOVA results for the Group and Condition main effects and interactions, as well as Group comparisons for the target and non-target conditions. Significance level is indicated as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Results are expressed as F-values (1,50).

3.2.1.1 ERT correlates with alcohol intake, performance, and attention

As shown in Figure 2, ERT power to targets in the right aIFC was positively correlated with response accuracy to targets ($r=0.29$, $p<0.05$) and with the total D2 attentional test score ($r=0.31$, $p<0.05$), indicating that right prefrontal theta power underlies successful target detection and higher attentional ability. In contrast, target-induced theta power estimated to the right aIFC was negatively correlated with the est-BAC ($r=-0.39$, $p<0.05$) and AUDIT scores ($r=-0.41$, $p>0.03$) in BDs. Correlations with the pIFC were very similar in pattern and significance indicating the association between lower theta power in the rIFC and higher levels of alcohol consumption which may underlie behavioral impairments.

[insert Figure 3.]

3.2.2 Co-oscillations in theta band (4-7 Hz) estimated with PLVs

To investigate functional interactions among the ROIs in real time, we estimated the degree of synchronous, in-phase co-oscillations by calculating the PLV in theta frequency band (Figure 3). The PLV between the right aIFC and pIFC areas to targets increased only in LDs $F(1,50)= 9.65$, $p<0.01$. Overall, synchronous co-oscillations were greater in LDs compared to BD as indicated by a main effect of group: $F(1,50)= 6.7$, $p<0.05$, and were greater for targets only, Group x Condition interaction: $F(1,50) = 6.84$, $p<0.05$. The synchrony was maximal in the time interval encompassing response execution (300-580 ms), indicating involvement of the ventrolateral prefrontal network. In addition, the PLV declined in BDs and was negatively correlated with the est-BAC ($r=-0.54$, $p<0.01$), suggesting that hazardous drinking levels are associated with dysregulation of network interactions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[insert Figure 4.]

4. DISCUSSION

The present study used a multimodal approach to examine theta band oscillatory activity and functional connectivity during target detection in adolescents emerging into adulthood as a function of binge drinking. The main findings can be summarized as follows: 1) the greatest event-related theta (ERT) was evoked by targets in the right inferior frontal cortex (rIFC); 2) It correlated with performance accuracy and selective attention scores, confirming the importance of rIFC in target detection, decision making, and response mapping. 3) Lower ERT to targets was observed in BDs compared to LDs in the rIFC. 4) Furthermore, ERT correlated negatively with the est-BAC and the AUDIT score in the BD group, lending support to the idea that lower attentional function in BDs is associated with heavy alcohol consumption. 5) Theta band co-oscillations between the right aIFC and pIFC were dysregulated in BDs indicating decreased connectivity of the prefrontal network during attentional engagement, response selection, and execution. The lower neural synchrony in BDs was associated with alcohol intake.

The IFC has been characterized as a critical substrate of cognitive control which refers to the ability to flexibly and strategically respond to contextual demands in agreement with past experiences and current goals (Aron et al., 2014; Brass et al., 2005; Chambers et al., 2009; Hampshire et al., 2010; Levy and Wagner, 2011; Munakata et al., 2011; Wiecki and Frank, 2013). Cognitive control is multiply determined by different underlying cognitive processes with specific contributions (Badre, 2008; Brass et al., 2005; Chikazoe, 2010; Criaud and Boulinguez, 2013; Koechlin et al., 2003; Levy and Wagner, 2011). It has been well established that the rIFC is engaged by tasks eliciting suppression of prepotent responses (Aron et al., 2014; Chikazoe, 2010; Garavan et al., 1999; Swick et al., 2011; Wessel and Aron, 2017). However, inherent to those paradigms are non-inhibitory confounds such as attention to salient, infrequent stimuli. Indeed, there is extensive evidence of the rIFC recruitment during attentional tasks that impose no or minimal inhibitory requirements (Erika-Florence et al., 2014; Hampshire et al., 2010; Hampshire and Sharp, 2015). Therefore, rIFC does not seem to solely specialize for motor inhibition but is activated by salient cues capturing attention (Hampshire et al., 2009, 2010). Studies using attentional manipulations confirm that the rIFC is an essential hub of the ventral attentional network (Chica et al., 2013; Corbetta et al., 2008; Corbetta and Shulman, 2002). Moreover, the ventral prefrontal network is engaged during deliberate target detection

1
2
3 (Hampshire et al., 2009; Indovina and Macaluso, 2007; Shulman et al., 2010; Vossel et al.,
4 2014) which relies on attentional control, response mapping, and maintenance of task sets
5
6 (Dosenbach et al., 2008). Results of the present study are aligned with such evidence as much
7
8 greater ERT was elicited by target stimuli in the rIFC due to its salience. In comparison,
9
10 equiprobable nontarget stimuli elicited virtually no theta in any of the frontal areas (Figure 1)
11
12 subsequent to evoking equivalent early activity in the visual cortex. This finding is consistent
13
14 with previous reports of greater theta to task-relevant stimuli with contributions of memory
15
16 retrieval, attention, working memory, decision making, and continuous monitoring (Cavanagh
17
18 and Frank, 2014; Cheyne et al., 2012; Halgren et al., 2015; Kovacevic et al., 2012; Nyhus and
19
20 Curran, 2010; Sauseng et al., 2010; Womelsdorf et al., 2010). Furthermore, theta power scales
21
22 with task difficulty and cognitive demands confirming its essential contributions to cognitive
23
24 control (Beaton et al., 2017; Ishii et al., 2014; Kovacevic et al., 2012; Rosen et al., 2016). In the
25
26 present study, ERT to targets in the rIFC correlated with better response accuracy and higher
27
28 scores on attentional test (Figure 2) which supports the involvement of theta in attentional
29
30 processing and responding to task-relevant stimuli.

31
32 Binge drinkers exhibited lower theta power to targets compared to LDs in the rIFC (Figure 1,
33
34 Table 3). Higher levels of alcohol consumption were associated with lower theta in BDs (Figure
35
36 2), suggesting that it may potentially serve as an index of reduced cognitive capacity in young,
37
38 highly functioning individuals as a function of their drinking levels. Group differences in task
39
40 performance eluded statistical significance although BDs showed a marginal tendency to
41
42 respond with longer reaction times and lower accuracy (Table 2). On a behavioral level, these
43
44 results are in line with the attentional deficits observed in BDs (Lannoy et al., 2017) and in AUD
45
46 individuals (Maurage et al., 2014; Muller-Oehring and Schulte, 2014; Oscar-Berman and
47
48 Marinković, 2007; Tedstone and Coyle, 2004), suggesting that impaired attentional control might
49
50 contribute to alcohol-related problems. Indeed, it has been well established that alcoholics
51
52 manifest reduced ERT power during cognitive tasks (Campanella et al., 2009; Kamarajan and
53
54 Porjesz, 2015; Pandey et al., 2012; Porjesz et al., 2005; Rangaswamy and Porjesz, 2014).
55
56 Previous EEG studies have examined ERT with equiprobable Go/NoGo tasks in individuals with
57
58 AUD and reported decreased theta power to both conditions (Kamarajan et al., 2004; Pandey et
59
60 al., 2016). Similarly, a recent EEG study that used the same paradigm in a shared sample as

1
2
3 the current study (López-Caneda et al., 2017) reported reduced power in delta and theta bands
4 in BDs to both target and non-target stimuli. These findings were interpreted as reflecting
5 dysfunctional response execution and response inhibition. They were partially replicated in our
6 study inasmuch as we found lower ERT in BDs. However, our results clearly indicate much
7 greater neural engagement during target (Go) trials suggesting that the observed activity was
8 primarily elicited by attention-inducing, target-relevant processes. In contrast, theta activity to
9 task irrelevant NoGo trials was barely above baseline levels indicating negligible contributions of
10 response inhibition. Furthermore, our method has permitted a more refined spatial specificity
11 indicating selective sensitivity of the rIFC to higher levels of alcohol consumption with lower ERT
12 to target trials in BDs. Additional studies are needed to explore these discrepancies, but they
13 may be due to different data analysis methods and may reflect better spatial sensitivity of the
14 MEG signal.

15
16
17 In a series of aMEG studies using different executive tasks and alcohol challenge in social
18 drinkers, we have observed attenuated ERT in the prefrontal cortex during alcohol intoxication
19 especially under higher cognitive load (Beaton et al., 2017; Kovacevic et al., 2012; Marinkovic et
20 al., 2012; Rosen et al., 2016). Lower theta has also been reported in high-risk offspring of
21 individuals with AUD (Kamarajan and Porjesz, 2015). Taken together, lower task-induced theta
22 oscillations have been proposed as an endophenotype for susceptibility to AUD (Andrew and
23 Fein, 2010; Hodgkinson et al., 2010; Porjesz et al., 2005; Rangaswamy and Porjesz, 2008;
24 Salvatore et al., 2015).

25
26
27 Target processing relies on sustained selective attention. Imaging studies confirm that the rIFC
28 may be involved in evaluating significance of salient events across sensory modalities (Downar
29 et al., 2002). Its sensitivity to visual salience (Vossel et al., 2011) is supported by evidence from
30 single-cell recordings in monkeys indicating that the IFC receives detailed visual information
31 from the ventral visual pathway (Sakagami and Pan, 2007; Sakagami and Tsutsui, 1999;
32 Scialidhe et al., 1999). Functional specialization and material-specific sensitivity of micro-
33 domains in the IFC has been shown with human intracranial recordings (Marinkovic et al.,
34 2000). Attentional processing is essential not only for detecting relevant stimuli but also for
35 managing distraction in a broader context of goal-dependent constraints. Altered behavioral and
36 brain-based indices of distractibility have been observed in abstinent alcoholics (Ahveninen et
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 al., 2000) while acute alcohol intoxication selectively impairs attention to novelty (Marinkovic et
4 al., 2001). Increased impulsivity and vulnerability to distraction are strongly implicated in alcohol
5 and drug addiction (Evenden, 1999). Inability to exert inhibitory control over drinking may result
6 from an interaction of genetic predisposition and drug-induced deficits (Begleiter and Porjesz,
7 1999; Dick et al., 2010). IFC has been implicated in top-down attentional control and emotional
8 reappraisal (Goldin et al., 2008; Green et al., 2011) and it may subserve the integration of
9 attentional control, decision making, and top-down regulative influences, and their degradation
10 in addiction (Goldstein and Volkow, 2002; Koob and Volkow, 2010).

11
12
13
14
15
16
17
18 Drawing on excellent temporal resolution of the MEG signal, the present study examined co-
19 oscillations between the two principal activated areas in the rIFC in real time (Figure 3). We
20 calculated PLVs in theta band to quantify consistency of the oscillatory phase between the aIFC
21 and pIFC (Lachaux et al., 1999). Co-oscillations are known to transiently increase at the time of
22 active interactions between two ROIs of interest (Beaton et al., 2017; Klopp et al., 2000; Tallon-
23 Baudry and Bertrand, 1999). Theta oscillations are particularly suitable to investigate cross-
24 cortical integration as they are primarily generated in superficial cortical layers (Halgren et al.,
25 2015, 2018) and are sensitive to episodic encoding and retrieval (Anderson et al., 2010;
26 Hanslmayr and Staudigl, 2014; Hasselmo and Stern, 2014; Klimesch et al., 2010; Womelsdorf
27 et al., 2010). As shown in Figure 3, co-oscillations between the aIFC and pIFC were negligible
28 on nontarget trials in both groups. In contrast, they were significantly greater to target stimuli in
29 LDs during the time of attentional selection, response preparation, and execution. This suggests
30 that the aIFC and pIFC were co-active during attentional control and motor response in LDs.
31 Given their functional synchronization, the two areas appear to be linked in an interactive IFC
32 network which is consistent with other evidence showing a high degree of interconnectivity in
33 the IFC (Goulas et al., 2012). However, due to potential confounds due to field spread
34 (Schoffelen and Gross, 2009), these findings should be considered with caution. The
35 converging findings suggest that the IFC may not be organized into separable, specialized
36 modules based on their specific functional profiles but that they are flexibly engaged and
37 integrated by adaptive contextual maintenance (Reynolds et al., 2012). Studies investigating the
38 functional organization of the IFC by manipulating hierarchical levels of rule representation have
39 confirmed a lack of spatial differentiation across different representational levels (Pischedda et
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 al., 2017). On that view, the contributions of the IFC to cognitive function are represented in
4 spatially overlapping areas that are fluidly engaged to conform to task rules. Other studies have
5 also suggested that the IFC and the neighboring anterior insular cortex represent a core hub for
6 rule processing and maintenance of task sets, integrating different dimensions of cognitive
7 control (Dosenbach et al., 2007; Menon and Uddin, 2010).
8
9

10
11
12 Our results indicate that the oscillatory synchrony within the IFC was dysregulated in BDs and
13 their PLV decrease was associated with alcohol intake (Figure 3). Previous studies reported that
14 alcohol intoxication abolished PLV during a task evoking cognitive control (Beaton et al., 2017;
15 Ehlers et al., 2012).. Altered connectivity in BDs has also been reported during wakeful resting-
16 state as reflected in PLV measured with MEG (Correas et al., 2015). Similarly, resting-state
17 functional connectivity MRI studies in individuals with AUD have found lower connectivity in the
18 executive control (Kim et al., 2017; Müller-Oehring et al., 2015; Weiland et al., 2014) and reward
19 networks (Camchong et al., 2013; Wang et al., 2016). Thus, the attenuated co-oscillations to
20 targets observed in the current study may indicate suboptimal integrative processing that could
21 underlie cognitive deficits in BDs.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

5. CONCLUSIONS AND LIMITATIONS

Taken together, these results confirm essential contributions of the rIFC to attentional control in a target detection task which relies on sustained and selective attention, response selection and execution, and performance monitoring. Lower ERT to target stimuli was observed in the rIFC in adolescents emerging into adulthood as a function of BD. It was associated with higher alcohol intake and worse performance accuracy and scores on a test of attention further indicating an impairment in attentional control in BDs. Heavy consumption was further associated with deficient co-oscillations in the rIFC that may underlie impaired cognitive functions in BDs. These indices may assist in devising interventions aimed at reducing drinking levels in young emerging adults.

Given the cross-sectional nature of this type of observational cohort studies, it is not possible to make causal inferences about the main factors underlying the loss of cognitive control. Although the observed correlations between the neural indices and alcohol intake suggest that binge drinking is associated with decrease cognitive control, it cannot be excluded that these differences precede alcohol consumption.

Declaration of interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This study was supported by the projects SPI/2010/134 and SPI/2010/051 from the Spanish Ministry of Health and Social Politics (National Plan of Drugs), and the National Institutes of Health, (R01-AA016624). Eduardo López-Caneda was supported by the Postdoctoral Fellowship of the Portuguese Foundation for Science and Technology SFRH/BPD/109750/2015.

Acknowledgement

We thank Pablo Cuesta and Burke Rosen for their technical assistance.

6. REFERENCES

- Ahveninen J, Jääskeläinen IP, Pekkonen E, et al. (2000) Increased distractibility by task-irrelevant sound changes in abstinent alcoholics. *Alcoholism, clinical and experimental research* 24(12): 1850–4.
- American Psychiatric Association (2000) Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR. Washington, DC: American Psychiatric Association.
- Anderson KL, Rajagovindan R, Ghacibeh GA, et al. (2010) Theta oscillations mediate interaction between prefrontal cortex and medial temporal lobe in human memory. *Cerebral cortex (New York, N.Y. : 1991)* 20(7): 1604–12. DOI: 10.1093/cercor/bhp223.
- Andrew C and Fein G (2010) Induced theta oscillations as biomarkers for alcoholism. *Clinical Neurophysiology* 121(3): 350–358. DOI: 10.1016/j.clinph.2009.11.080.
- Aron AR, Robbins TW and Poldrack RA (2014) Inhibition and the right inferior frontal cortex: one decade on. *Trends in Cognitive Sciences* 18(4): 177–185. DOI: 10.1016/j.tics.2013.12.003.
- Badre D (2008) Cognitive control, hierarchy, and the rostro-caudal organization of the frontal lobes. *Trends in Cognitive Sciences* 12(5): 193–200. DOI: 10.1016/j.tics.2008.02.004.
- Beaton LE, Azma S and Marinkovic K (2017) When the brain changes its mind: Oscillatory dynamics of conflict processing and response switching in a flanker task during alcohol challenge. *PLoS one*.
- Bechara A, Damasio AR, Damasio H, et al. (1994) Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* 50(1–3): 7–15.
- Becoña E, Fernández del Río E, López A, et al. (2011) [The Short Nicotine Dependence Syndrome Scale (NDSS-S) in Spanish smokers]. *Psicothema* 23(1): 126–32.
- Begleiter H and Porjesz B (1999) What is inherited in the predisposition toward alcoholism? A proposed model. *Alcoholism, clinical and experimental research* 23(7): 1125–35.
- Brass M, Derrfuss J, Forstmann BU, et al. (2005) The role of the inferior frontal junction area in cognitive control. *Trends in Cognitive Sciences* 9(7): 314–316. DOI: 10.1016/j.tics.2005.05.001.
- Brickenkamp R and Seisdodos N (2009) D2, test de atención. [D2 Attention test] (3rd ed.). TEA, Madrid, Spain.
- Brier MR, Ferree TC, Maguire MJ, et al. (2010) Frontal theta and alpha power and coherence changes are modulated by semantic complexity in Go/NoGo tasks. *International journal of psychophysiology : official journal of the International Organization of Psychophysiology* 78(3): 215–24. DOI: 10.1016/j.ijpsycho.2010.07.011.
- Camchong J, Stenger VA and Fein G (2013) Resting-state synchrony in short-term versus long-term abstinent alcoholics. *Alcoholism, clinical and experimental research* 37(5): 794–803. DOI: 10.1111/acer.12037.
- Campanella S, Petit G, Maurage P, et al. (2009) Chronic alcoholism: insights from neurophysiology. *Neurophysiologie clinique = Clinical neurophysiology* 39(4–5): 191–207. DOI: 10.1016/j.neucli.2009.08.002.
- Casey BJ, Tottenham N, Liston C, et al. (2005) Imaging the developing brain: what have we learned about cognitive development? *Trends in Cognitive Sciences* 9(3): 104–110. DOI: 10.1016/j.tics.2005.01.011.

- 1
2
3 Cavanagh JF and Frank MJ (2014) Frontal theta as a mechanism for cognitive control. *Trends*
4 *in Cognitive Sciences* 18(8): 414–421. DOI: 10.1016/j.tics.2014.04.012.
- 5 Chambers CD, Garavan H and Bellgrove M a (2009) Insights into the neural basis of response
6 inhibition from cognitive and clinical neuroscience. *Neuroscience and biobehavioral*
7 *reviews* 33(5): 631–46. DOI: 10.1016/j.neubiorev.2008.08.016.
- 8
9 Cheyne DO, Ferrari P and Cheyne JA (2012) Intended actions and unexpected outcomes:
10 automatic and controlled processing in a rapid motor task. *Frontiers in Human*
11 *Neuroscience* 6: 237. DOI: 10.3389/fnhum.2012.00237.
- 12 Chica AB, Bartolomeo P and Lupiáñez J (2013) Two cognitive and neural systems for
13 endogenous and exogenous spatial attention. *Behavioural brain research* 237: 107–23.
14 DOI: 10.1016/j.bbr.2012.09.027.
- 15 Chikazoe J (2010) Localizing performance of go/no-go tasks to prefrontal cortical subregions.
16 *Current Opinion in Psychiatry* 23(3): 267–272. DOI: 10.1097/YCO.0b013e3283387a9f.
- 17
18 Colrain IM, Sullivan E V, Ford JM, et al. (2011) Frontally mediated inhibitory processing and
19 white matter microstructure: age and alcoholism effects. *Psychopharmacology* 213(4):
20 669–79. DOI: 10.1007/s00213-010-2073-7.
- 21
22 Corbetta M and Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the
23 brain. *Nature reviews. Neuroscience* 3(3): 201–15. DOI: 10.1038/nm755.
- 24 Corbetta M, Patel G and Shulman GL (2008) The reorienting system of the human brain: from
25 environment to theory of mind. *Neuron* 58(3): 306–24. DOI: 10.1016/j.neuron.2008.04.017.
- 26
27 Correas A, Rodriguez Holguín S, Cuesta P, et al. (2015) Exploratory Analysis of Power
28 Spectrum and Functional Connectivity During Resting State in Young Binge Drinkers: A
29 MEG Study. *International Journal of Neural Systems* 25(03): 1550008. DOI:
30 10.1142/S0129065715500082.
- 31 Correas A, Cuesta P, López-Caneda E, et al. (2016) Functional and structural brain connectivity
32 of young binge drinkers: A follow-up study. *Scientific Reports* 6. DOI: 10.1038/srep31293.
- 33
34 Courtney KE and Polich J (2009) Binge Drinking in Young Adults: Data, Definitions and
35 Determinants. *Psychol bull* 135(1): 142–156. DOI: 10.1037/a0014414.Binge.
- 36
37 Crabbe JC, Harris RA and Koob GF (2011) Preclinical studies of alcohol binge drinking. *Annals*
38 *of the New York Academy of Sciences* 1216(1): 24–40. DOI: 10.1111/j.1749-
39 6632.2010.05895.x.
- 40 Crews FT and Boettiger CA (2009) Impulsivity, frontal lobes and risk for addiction.
41 *Pharmacology, biochemistry, and behavior* 93(3). Elsevier Inc.: 237–47. DOI:
42 10.1016/j.pbb.2009.04.018.
- 43
44 Crews FT, Vetreno RP, Broadwater MA, et al. (2016) Adolescent Alcohol Exposure Persistently
45 Impacts Adult Neurobiology and Behavior. *Pharmacological reviews* 68(4): 1074–1109.
46 DOI: 10.1124/pr.115.012138.
- 47 Criaud M and Boulinguez P (2013) Have we been asking the right questions when assessing
48 response inhibition in go/no-go tasks with fMRI? A meta-analysis and critical review.
49 *Neuroscience and biobehavioral reviews* 37(1): 11–23. DOI:
50 10.1016/j.neubiorev.2012.11.003.
- 51 Cservenka A and Brumback T (2017) The Burden of Binge and Heavy Drinking on the Brain:
52 Effects on Adolescent and Young Adult Neural Structure and Function. *Frontiers in*
53 *psychology* 8: 1111. DOI: 10.3389/fpsyg.2017.01111.
- 54
55 Dale AM, Liu AK, Fischl BR, et al. (2000) Dynamic statistical parametric mapping: combining
56 fMRI and MEG for high-resolution imaging of cortical activity. *Neuron* 26(1): 55–67.
- 57
58
59
60

- 1
2
3 de Wit H (2009) Impulsivity as a determinant and consequence of drug use: a review of
4 underlying processes. *Addiction biology* 14(1): 22–31. DOI: 10.1111/j.1369-
5 1600.2008.00129.x.
- 6 Delorme A and Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial
7 EEG dynamics including independent component analysis. *Journal of neuroscience*
8 *methods* 134(1): 9–21. DOI: 10.1016/j.jneumeth.2003.10.009.
- 9
10 Derogatis LR and Melisaratos N (1983) The Brief Symptom Inventory: an introductory report.
11 *Psychological medicine* 13(3): 595–605.
- 12 Diamond A (2013) Executive functions. *Annual review of psychology* 64(1): 135–68. DOI:
13 10.1146/annurev-psych-113011-143750.
- 14
15 Dick DM, Smith G, Olausson P, et al. (2010) Understanding the construct of impulsivity and its
16 relationship to alcohol use disorders. *Addiction Biology* 15(2): 217–226. DOI:
17 10.1111/j.1369-1600.2009.00190.x.
- 18 Dosenbach NUF, Fair DA, Miezin FM, et al. (2007) Distinct brain networks for adaptive and
19 stable task control in humans. *Proceedings of the National Academy of Sciences* 104(26):
20 11073–11078. DOI: 10.1073/pnas.0704320104.
- 21
22 Dosenbach NUF, Fair D a, Cohen AL, et al. (2008) A dual-networks architecture of top-down
23 control. *Trends in cognitive sciences* 12(3): 99–105. DOI: 10.1016/j.tics.2008.01.001.
- 24
25 Downar J, Crawley AP, Mikulis DJ, et al. (2002) A cortical network sensitive to stimulus salience
26 in a neutral behavioral context across multiple sensory modalities. *Journal of*
27 *neurophysiology* 87(1): 615–20. DOI: 10.1152/jn.00636.2001.
- 28 Ehlers CL, Wills DN and Havstad J (2012) Ethanol reduces the phase locking of neural activity
29 in human and rodent brain. *Brain research* 1450: 67–79. DOI:
30 10.1016/j.brainres.2012.02.039.
- 31
32 Erika-Florence M, Leech R and Hampshire A (2014) A functional network perspective on
33 response inhibition and attentional control. *Nature Communications* 5: 4073. DOI:
34 10.1038/ncomms5073.
- 35 Eurobarometer (2010) EU citizens' attitudes towards alcohol. Special Eurobarometer 331/Wave
36 72.3.
- 37
38 Evenden JL (1999) Varieties of impulsivity. *Psychopharmacology* 146(4): 348–61.
- 39
40 Fan J, McCandliss BD, Fossella J, et al. (2005) The activation of attentional networks.
41 *Neuroimage* 26(2): 471–9. DOI: 10.1016/j.neuroimage.2005.02.004.
- 42
43 Fischl B, Sereno MI, Tootell RB, et al. (1999) High-resolution intersubject averaging and a
44 coordinate system for the cortical surface. *Human brain mapping* 8(4): 272–84.
- 45
46 Figueira-Ares R, Cadaveira F, Rodríguez Holguín S, et al. (2017) Electrophysiological
47 Anomalies in Face-Name Memory Encoding in Young Binge Drinkers. *Frontiers in*
48 *psychiatry* 8: 216. DOI: 10.3389/fpsy.2017.00216.
- 49
50 Garavan H, Ross TJ and Stein E a (1999) Right hemispheric dominance of inhibitory control: an
51 event-related functional MRI study. *Proceedings of the National Academy of Sciences of*
52 *the United States of America* 96(14): 8301–6.
- 53
54 Gitelman DR, Nobre AC, Parrish TB, et al. (1999) A large-scale distributed network for covert
55 spatial attention: further anatomical delineation based on stringent behavioural and
56 cognitive controls. *Brain : a journal of neurology* 122 (Pt 6): 1093–106.
- 57
58 Gogtay N, Giedd JN, Lusk L, et al. (2004) Dynamic mapping of human cortical development
59 during childhood through early adulthood. *Proceedings of the National Academy of*
60 *Sciences of the United States of America* 101(21): 8174–9. DOI:

1
2
3 10.1073/pnas.0402680101.

4 Goldin PR, McRae K, Ramel W, et al. (2008) The Neural Bases of Emotion Regulation:
5 Reappraisal and Suppression of Negative Emotion. *Biological Psychiatry* 63(6): 577–586.
6 DOI: 10.1016/j.biopsych.2007.05.031.

7
8 Goldstein RZ and Volkow ND (2002) Drug Addiction and Its Underlying Neurobiological Basis:
9 Neuroimaging Evidence for the Involvement of the Frontal Cortex. *American Journal of*
10 *Psychiatry* 159(10): 1642–1652. DOI: 10.1176/appi.ajp.159.10.1642.

11 Goulas A, Uylings HBM and Stiers P (2012) Unravelling the Intrinsic Functional Organization of
12 the Human Lateral Frontal Cortex: A Parcellation Scheme Based on Resting State fMRI.
13 *Journal of Neuroscience* 32(30): 10238–10252. DOI: 10.1523/JNEUROSCI.5852-11.2012.

14
15 Gramfort A, Luessi M, Larson E, et al. (2014) MNE software for processing MEG and EEG data.
16 *NeuroImage* 86: 446–60. DOI: 10.1016/j.neuroimage.2013.10.027.

17 Green JJ, Doesburg SM, Ward LM, et al. (2011) Electrical Neuroimaging of Voluntary
18 Audiospatial Attention: Evidence for a Supramodal Attention Control Network. *Journal of*
19 *Neuroscience* 31(10): 3560–3564. DOI: 10.1523/JNEUROSCI.5758-10.2011.

20
21 Guillamón M, Solé A and Farran J (1999) Test para la identificación de trastornos por uso de
22 alcohol (AUDIT): Traducción y validación del AUDIT al catalán y el castellano. *Adicciones*
23 11: 337–347.

24 Halgren E, Kaestner E, Marinkovic K, et al. (2015) Laminar profile of spontaneous and evoked
25 theta: Rhythmic modulation of cortical processing during word integration.
26 *Neuropsychologia* 76: 108–24. DOI: 10.1016/j.neuropsychologia.2015.03.021.

27
28 Halgren M, Fabó D, Ulbert I, et al. (2018) Superficial Slow Rhythms Integrate Cortical
29 Processing in Humans. *Scientific Reports* 8(1): 2055. DOI: 10.1038/s41598-018-20662-0.

30 Hampshire A and Sharp D (2015) Inferior PFC Subregions Have Broad Cognitive Roles. *Trends*
31 *in Cognitive Sciences* 19(12): 712–713. DOI: 10.1016/j.tics.2015.09.010.

32
33 Hampshire A, Thompson R, Duncan J, et al. (2009) Selective tuning of the right inferior frontal
34 gyrus during target detection. *Cognitive, affective & behavioral neuroscience* 9(1): 103–12.
35 DOI: 10.3758/CABN.9.1.103.

36 Hampshire A, Chamberlain SR, Monti MM, et al. (2010) The role of the right inferior frontal
37 gyrus: inhibition and attentional control. *NeuroImage* 50(3): 1313–1319. DOI:
38 10.1016/j.neuroimage.2009.12.109.

39
40 Hanslmayr S and Staudigl T (2014) How brain oscillations form memories — A processing
41 based perspective on oscillatory subsequent memory effects. *NeuroImage* 85: 648–655.
42 DOI: 10.1016/j.neuroimage.2013.05.121.

43 Hanslmayr S, Pastötter B, Bäuml K-H, et al. (2008) The electrophysiological dynamics of
44 interference during the Stroop task. *Journal of cognitive neuroscience* 20(2): 215–25. DOI:
45 10.1162/jocn.2008.20020.

46
47 Hasselmo ME and Stern CE (2014) Theta rhythm and the encoding and retrieval of space and
48 time. *NeuroImage* 85: 656–666. DOI: 10.1016/j.neuroimage.2013.06.022.

49 Hermens DF, Lagopoulos J, Tobias-Webb J, et al. (2013) Pathways to alcohol-induced brain
50 impairment in young people: A review. *Cortex* 49(1): 3–17.

51
52 Hingson RW and White A (2013) Trends in Extreme Binge Drinking Among US High School
53 Seniors. *JAMA Pediatrics* 167(11): 996. DOI: 10.1001/jamapediatrics.2013.3083.

54
55 Hingson RW, Zha W and White AM (2017) Drinking Beyond the Binge Threshold: Predictors,
56 Consequences, and Changes in the U.S. *American journal of preventive medicine* 52(6):
57 717–727. DOI: 10.1016/j.amepre.2017.02.014.

- 1
2
3 Hodgkinson CA, Enoch M-A, Srivastava V, et al. (2010) Genome-wide association identifies
4 candidate genes that influence the human electroencephalogram. *Proceedings of the*
5 *National Academy of Sciences of the United States of America* 107(19): 8695–700. DOI:
6 10.1073/pnas.0908134107.
- 7 Indovina I and Macaluso E (2007) Dissociation of stimulus relevance and saliency factors during
8 shifts of visuospatial attention. *Cerebral cortex (New York, N.Y. : 1991)* 17(7): 1701–11.
9 DOI: 10.1093/cercor/bhl081.
- 10 Ishii R, Canuet L, Ishihara T, et al. (2014) Frontal midline theta rhythm and gamma power
11 changes during focused attention on mental calculation: an MEG beamformer analysis.
12 *Frontiers in human neuroscience* 8: 406. DOI: 10.3389/fnhum.2014.00406.
- 13
14 Jacobus J and Tapert SF (2013) Neurotoxic effects of alcohol in adolescence. *Annual review of*
15 *clinical psychology* 9: 703–21. DOI: 10.1146/annurev-clinpsy-050212-185610.
- 16
17 Jensen O and Tesche CD (2002) Frontal theta activity in humans increases with memory load in
18 a working memory task. *The European journal of neuroscience* 15(8): 1395–9.
- 19
20 Johnston L (2011) Monitoring the Future: National Results on Adolescent Drug Use: Overview
21 of Key Findings.
- 22
23 Kamarajan C and Porjesz B (2015) Advances in Electrophysiological Research. *Alcohol*
24 *research : current reviews* 37(1): 53–87.
- 25
26 Kamarajan C, Porjesz B, Jones KA, et al. (2004) The role of brain oscillations as functional
27 correlates of cognitive systems : a study of frontal inhibitory control in alcoholism. 51: 155–
28 180. DOI: 10.1016/j.ijpsycho.2003.09.004.
- 29
30 Kim S, Im S, Lee J, et al. (2017) Disrupted Control Network Connectivity in Abstinent Patients
31 with Alcohol Dependence. *Psychiatry Investigation* 14(3): 325. DOI:
32 10.4306/pi.2017.14.3.325.
- 33
34 Klimesch W, Freunberger R and Sauseng P (2010) Oscillatory mechanisms of process binding
35 in memory. *Neuroscience & Biobehavioral Reviews* 34(7): 1002–1014. DOI:
36 10.1016/j.neubiorev.2009.10.004.
- 37
38 Klinke ME, Hjaltason H, Tryggvadóttir GBFR, et al. (2017) Hemispatial neglect following right
39 hemisphere stroke: clinical course and sensitivity of diagnostic tasks. *Topics in stroke*
40 *rehabilitation*: 1–11. DOI: 10.1080/10749357.2017.1394632.
- 41
42 Klopp J, Marinkovic K, Chauvel P, et al. (2000) Early widespread cortical distribution of coherent
43 fusiform face selective activity. *Human brain mapping* 11(4): 286–93.
- 44
45 Koechlin E, Ody C and Kouneiher F (2003) The Architecture of Cognitive Control in the Human
46 Prefrontal Cortex. *Science* 302(5648): 1181–1185. DOI: 10.1126/science.1088545.
- 47
48 Koob GF and Volkow ND (2010) Neurocircuitry of addiction. *Neuropsychopharmacology : official*
49 *publication of the American College of Neuropsychopharmacology* 35(1). American
50 College of Neuropsychopharmacology: 217–38. DOI: 10.1038/npp.2009.110.
- 51
52 Kovacevic S, Azma S, Irimia A, et al. (2012) Theta oscillations are sensitive to both early and
53 late conflict processing stages: Effects of alcohol intoxication. *PLoS ONE* 7(8): 1–11. DOI:
54 10.1371/journal.pone.0043957.
- 55
56 Lachaux JP, Rodriguez E, Martinerie J, et al. (1999) Measuring phase synchrony in brain
57 signals. *Human brain mapping* 8(4): 194–208.
- 58
59 Lannoy S, Heeren A, Moyaerts N, et al. (2017) Differential impairments across attentional
60 networks in binge drinking. *Psychopharmacology* 234(7): 1059–1068. DOI:
10.1007/s00213-017-4538-4.
- Lebel C and Beaulieu C (2011) Longitudinal development of human brain wiring continues from

- 1
2
3 childhood into adulthood. *The Journal of neuroscience : the official journal of the Society*
4 *for Neuroscience* 31(30): 10937–47. DOI: 10.1523/JNEUROSCI.5302-10.2011.
- 5 Levy BJ and Wagner AD (2011) Cognitive control and right ventrolateral prefrontal cortex:
6 reflexive reorienting, motor inhibition, and action updating. *Annals of the New York*
7 *Academy of Sciences* 1224(1): 40–62. DOI: 10.1111/j.1749-6632.2011.05958.x.
- 8
9 Lin F-H, Witzel T, Hämäläinen MS, et al. (2004) Spectral spatiotemporal imaging of cortical
10 oscillations and interactions in the human brain. *NeuroImage* 23(2): 582–95. DOI:
11 10.1016/j.neuroimage.2004.04.027.
- 12 López-Caneda E, Cadaveira F, Crego A, et al. (2012) Hyperactivation of right inferior frontal
13 cortex in young binge drinkers during response inhibition: a follow-up study. *Addiction*
14 *(Abingdon, England)* 107(10): 1796–808. DOI: 10.1111/j.1360-0443.2012.03908.x.
- 15 López-Caneda E, Cadaveira F, Crego A, et al. (2013) Effects of a persistent binge drinking
16 pattern of alcohol consumption in young people: a follow-up study using event-related
17 potentials. *Alcohol and alcoholism (Oxford, Oxfordshire)* 48(4): 464–71. DOI:
18 10.1093/alcalc/agt046.
- 19
20 López-Caneda E, Rodríguez Holguín S, Correas Á, et al. (2017) Binge drinking affects brain
21 oscillations linked to motor inhibition and execution. *Journal of Psychopharmacology*
22 31(7). DOI: 10.1177/0269881116689258.
- 23 Marinkovic K, Trebon P, Chauvel P, et al. (2000) Localised face processing by the human
24 prefrontal cortex: face-selective intracerebral potentials and post-lesion deficits. *Cognitive*
25 *neuropsychology* 17(1): 187–99. DOI: 10.1080/026432900380562.
- 26
27 Marinkovic K, Halgren E and Maltzman I (2001) Arousal-related P3a to novel auditory stimuli is
28 abolished by a moderately low alcohol dose. *Alcohol and alcoholism (Oxford, Oxfordshire)*
29 36(6): 529–39.
- 30 Marinkovic K, Dhond RP, Dale AM, et al. (2003) Spatiotemporal dynamics of modality-specific
31 and supramodal word processing. *Neuron* 38(3): 487–97.
- 32
33 Marinkovic K, Rosen BQ, Cox B, et al. (2012) Event-Related Theta Power during Lexical-
34 Semantic Retrieval and Decision Conflict is Modulated by Alcohol Intoxication:
35 Anatomically Constrained MEG. *Frontiers in Psychology* 3: 121. DOI:
36 10.3389/fpsyg.2012.00121.
- 37 Maris E and Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data.
38 *Journal of Neuroscience Methods* 164(1): 177–190.
- 39
40 Maurage P, de Timary P, Billieux J, et al. (2014) Attentional alterations in alcohol dependence
41 are underpinned by specific executive control deficits. *Alcoholism, clinical and*
42 *experimental research* 38(7): 2105–12. DOI: 10.1111/acer.12444.
- 43 Menon V and Uddin LQ (2010) Saliency, switching, attention and control: a network model of
44 insula function. *Brain Structure and Function* 214(5–6): 655–667. DOI: 10.1007/s00429-
45 010-0262-0.
- 46
47 Muller-Oehring EM and Schulte T (2014) Alcohol and the Nervous System, Cognition emotion
48 and attention. In: Sullivan E V and Pfefferbaum A (eds) *Handbook of Clinical Neurology*,
49 pp. 341–349.
- 50 Müller-Oehring EM, Jung Y-C, Pfefferbaum A, et al. (2015) The Resting Brain of Alcoholics.
51 *Cerebral cortex (New York, N.Y. : 1991)* 25(11): 4155–68. DOI: 10.1093/cercor/bhu134.
- 52
53 Munakata Y, Herd SA, Chatham CH, et al. (2011) A unified framework for inhibitory control.
54 *Trends in cognitive sciences* 15(10): 453–9. DOI: 10.1016/j.tics.2011.07.011.
- 55
56 Naimi TS, Nelson DE and Brewer RD (2010) The Intensity of Binge Alcohol Consumption
57
58
59
60

- 1
2
3 Among U.S. Adults. *American Journal of Preventive Medicine* 38(2): 201–207. DOI:
4 10.1016/j.amepre.2009.09.039.
- 5 National Institute of Alcohol and Alcoholism (2004) NIAAA council approves definition of binge
6 drinking. *NIAAA newsletter*, 3 p. 3.
- 7
8 National Institute on Alcohol Abuse and Alcoholism (n.d.) Drinking Levels Defined | National
9 Institute on Alcohol Abuse and Alcoholism (NIAAA). Available at:
10 [https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-](https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking)
11 [drinking](https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking) (accessed 18 December 2017).
- 12 Nobre AC, Sebestyen GN, Gitelman DR, et al. (1997) Functional localization of the system for
13 visuospatial attention using positron emission tomography. *Brain : a journal of neurology*
14 120 (Pt 3): 515–33.
- 15 Nyhus E and Curran T (2010) Functional role of gamma and theta oscillations in episodic
16 memory. *Neuroscience & Biobehavioral Reviews* 34(7): 1023–1035. DOI:
17 10.1016/j.neubiorev.2009.12.014.
- 18
19 Oostenveld R, Fries P, Maris E, et al. (2011) FieldTrip: Open source software for advanced
20 analysis of MEG, EEG, and invasive electrophysiological data. *Computational intelligence*
21 *and neuroscience* 2011: 156869. DOI: 10.1155/2011/156869.
- 22
23 Oscar-Berman M and Marinković K (2007) Alcohol: effects on neurobehavioral functions and the
24 brain. *Neuropsychology review* 17(3): 239–57. DOI: 10.1007/s11065-007-9038-6.
- 25 Pandey AK, Kamarajan C, Rangaswamy M, et al. (2012) Event-Related Oscillations in
26 Alcoholism Research: A Review. *Journal of addiction research & therapy* Suppl 7(1). NIH
27 Public Access. DOI: 10.4172/2155-6105.S7-001.
- 28
29 Pandey AK, Kamarajan C, Manz N, et al. (2016) Delta, theta, and alpha event-related
30 oscillations in alcoholics during Go/NoGo task: Neurocognitive deficits in execution,
31 inhibition, and attention processing. *Progress in Neuro-Psychopharmacology and*
32 *Biological Psychiatry* 65(February): 158–171. DOI: 10.1016/j.pnpbp.2015.10.002.
- 33 Patrick ME and Terry-McElrath YM (2017) High-intensity drinking by underage young adults in
34 the United States. *Addiction (Abingdon, England)* 112(1): 82–93. DOI: 10.1111/add.13556.
- 35
36 Patton JH, Stanford MS and Barratt ES (1995) Factor structure of the Barratt impulsiveness
37 scale. *Journal of clinical psychology* 51(6): 768–74.
- 38
39 Petit G, Maurage P, Kornreich C, et al. (2014) Binge drinking in adolescents: a review of
40 neurophysiological and neuroimaging research. *Alcohol and alcoholism (Oxford,*
41 *Oxfordshire)* 49(2): 198–206. DOI: 10.1093/alcalc/agt172.
- 42 Pischedda D, Görden K, Haynes J-D, et al. (2017) Neural Representations of Hierarchical Rule
43 Sets: The Human Control System Represents Rules Irrespective of the Hierarchical Level
44 to Which They Belong. *The Journal of Neuroscience* 37(50): 12281–12296. DOI:
45 10.1523/JNEUROSCI.3088-16.2017.
- 46 Porjesz B, Rangaswamy M, Kamarajan C, et al. (2005) The utility of neurophysiological markers
47 in the study of alcoholism. *Clinical neurophysiology : official journal of the International*
48 *Federation of Clinical Neurophysiology* 116(5): 993–1018. DOI:
49 10.1016/j.clinph.2004.12.016.
- 50
51 Rangaswamy M and Porjesz B (2008) Uncovering genes for cognitive (dys)function and
52 predisposition for alcoholism spectrum disorders: a review of human brain oscillations as
53 effective endophenotypes. *Brain research* 1235: 153–71. DOI:
54 10.1016/j.brainres.2008.06.053.
- 55
56 Rangaswamy M and Porjesz B (2014) Understanding alcohol use disorders with
57 neuroelectrophysiology. *Handbook of clinical neurology* 125: 383–414. DOI:
58
59
60

1
2
3 10.1016/B978-0-444-62619-6.00023-9.

4 Reynolds JR, O'Reilly RC, Cohen JD, et al. (2012) The function and organization of lateral
5 prefrontal cortex: a test of competing hypotheses. Gilbert S (ed.) *PloS one* 7(2): e30284.
6 DOI: 10.1371/journal.pone.0030284.

7
8 Rosen BQ, Padovan N and Marinkovic K (2016) Alcohol Hits You When It Is Hard: Intoxication,
9 Task Difficulty, and Theta Brain Oscillations. *Alcoholism: Clinical and Experimental*
10 *Research* 40(4): 743–752. DOI: 10.1111/acer.13014.

11 Sakagami M and Pan X (2007) Functional role of the ventrolateral prefrontal cortex in decision
12 making. *Current opinion in neurobiology* 17(2): 228–33. DOI: 10.1016/j.conb.2007.02.008.

13
14 Sakagami M and Tsutsui K (1999) The hierarchical organization of decision making in the
15 primate prefrontal cortex. *Neuroscience research* 34(2): 79–89.

16 Salvatore JE, Gottesman II and Dick DM (2015) Endophenotypes for Alcohol Use Disorder: An
17 Update on the Field. *Current addiction reports* 2(1): 76–90. DOI: 10.1007/s40429-015-
18 0046-y.

19
20 SAMHSA (2013) Substance Abuse and Mental Health Services Administration. *Results from the*
21 *2012 National Survey on Drug Use and Health*: 13–4795.

22 Sauseng P, Griesmayr B, Freunberger R, et al. (2010) Control mechanisms in working memory:
23 a possible function of EEG theta oscillations. *Neuroscience and biobehavioral reviews*
24 34(7): 1015–22. DOI: 10.1016/j.neubiorev.2009.12.006.

25
26 Scalaidhe SP, Wilson FA and Goldman-Rakic PS (1999) Face-selective neurons during passive
27 viewing and working memory performance of rhesus monkeys: evidence for intrinsic
28 specialization of neuronal coding. *Cerebral cortex (New York, N. Y. : 1991)* 9(5): 459–75.

29 Schoffelen J-M and Gross J (2009) Source connectivity analysis with MEG and EEG. *Human*
30 *Brain Mapping* 30(6): 1857–1865. DOI: 10.1002/hbm.20745.

31
32 Sherman LE, Rudie JD, Pfeifer JH, et al. (2014) Development of the default mode and central
33 executive networks across early adolescence: a longitudinal study. *Developmental*
34 *cognitive neuroscience* 10: 148–59. DOI: 10.1016/j.dcn.2014.08.002.

35 Shulman GL, Pope DLW, Astafiev S V, et al. (2010) Right hemisphere dominance during spatial
36 selective attention and target detection occurs outside the dorsal frontoparietal network.
37 *The Journal of neuroscience : the official journal of the Society for Neuroscience* 30(10):
38 3640–51. DOI: 10.1523/JNEUROSCI.4085-09.2010.

39
40 Silveri MM, Dager AD, Cohen-Gilbert JE, et al. (2016) Neurobiological signatures associated
41 with alcohol and drug use in the human adolescent brain. *Neuroscience and biobehavioral*
42 *reviews* 70: 244–259. DOI: 10.1016/j.neubiorev.2016.06.042.

43 Simmonds DJ, Hallquist MN, Asato M, et al. (2014) Developmental stages and sex differences
44 of white matter and behavioral development through adolescence: a longitudinal diffusion
45 tensor imaging (DTI) study. *NeuroImage* 92: 356–68. DOI:
46 10.1016/j.neuroimage.2013.12.044.

47
48 Squeglia LM, Jacobus J and Tapert SF (2014) The effect of alcohol use on human adolescent
49 brain structures and systems. In: *Handbook of clinical neurology*, pp. 501–510. DOI:
50 10.1016/B978-0-444-62619-6.00028-8.

51 Stephens DN and Duka T (2008) Review. Cognitive and emotional consequences of binge
52 drinking: role of amygdala and prefrontal cortex. *Philosophical transactions of the Royal*
53 *Society of London. Series B, Biological sciences* 363(1507): 3169–79. DOI:
54 10.1098/rstb.2008.0097.

55
56 Sullivan E V and Pfefferbaum A (2005) Neurocircuitry in alcoholism: a substrate of disruption
57
58
59
60

- and repair. *Psychopharmacology* 180(4): 583–94. DOI: 10.1007/s00213-005-2267-6.
- Swick D, Ashley V and Turken U (2011) Are the neural correlates of stopping and not going identical? Quantitative meta-analysis of two response inhibition tasks. *NeuroImage* 56(3): 1655–65. DOI: 10.1016/j.neuroimage.2011.02.070.
- Tallon-Baudry and Bertrand (1999) Oscillatory gamma activity in humans and its role in object representation. *Trends in cognitive sciences* 3(4): 151–162.
- Tedstone D and Coyle K (2004) Cognitive impairments in sober alcoholics: performance on selective and divided attention tasks. *Drug and alcohol dependence* 75(3): 277–86. DOI: 10.1016/j.drugalcdep.2004.03.005.
- Terry-McElrath YM and Patrick ME (2016) Intoxication and binge and high-intensity drinking among US young adults in their mid-20s. *Substance Abuse* 37(4): 597–605. DOI: 10.1080/08897077.2016.1178681.
- Varela F, Lachaux JP, Rodriguez E, et al. (2001) The brainweb: phase synchronization and large-scale integration. *Nature reviews. Neuroscience* 2(4): 229–39. DOI: 10.1038/35067550.
- Vossel S, Weidner R and Fink GR (2011) Dynamic Coding of Events within the Inferior Frontal Gyrus in a Probabilistic Selective Attention Task. *Journal of Cognitive Neuroscience* 23(2): 414–424. DOI: 10.1162/jocn.2010.21441.
- Vossel S, Geng JJ and Fink GR (2014) Dorsal and ventral attention systems: distinct neural circuits but collaborative roles. *The Neuroscientist: a review journal bringing neurobiology, neurology and psychiatry* 20(2): 150–9. DOI: 10.1177/1073858413494269.
- Walhovd KB, Fjell AM, Brown TT, et al. (2012) Long-term influence of normal variation in neonatal characteristics on human brain development. *Proceedings of the National Academy of Sciences of the United States of America* 109(49): 20089–94. DOI: 10.1073/pnas.1208180109.
- Wang C, Ulbert I, Schomer DL, et al. (2005) Responses of human anterior cingulate cortex microdomains to error detection, conflict monitoring, stimulus-response mapping, familiarity, and orienting. *The Journal of neuroscience: the official journal of the Society for Neuroscience* 25(3): 604–13. DOI: 10.1523/JNEUROSCI.4151-04.2005.
- Wang J, Fan Y, Dong Yue, et al. (2016) Alterations in Brain Structure and Functional Connectivity in Alcohol Dependent Patients and Possible Association with Impulsivity. Zuo X-N (ed.) *PLOS ONE* 11(8): e0161956. DOI: 10.1371/journal.pone.0161956.
- Weiland BJ, Sabbineni A, Calhoun VD, et al. (2014) Reduced left executive control network functional connectivity is associated with alcohol use disorders. *Alcoholism, clinical and experimental research* 38(9): 2445–53. DOI: 10.1111/acer.12505.
- Wessel JR and Aron AR (2017) On the Globality of Motor Suppression: Unexpected Events and Their Influence on Behavior and Cognition. *Neuron* 93(2): 259–280. DOI: 10.1016/j.neuron.2016.12.013.
- Wiecki T V and Frank MJ (2013) A computational model of inhibitory control in frontal cortex and basal ganglia. *Psychological review* 120(2): 329–55. DOI: 10.1037/a0031542.
- Windmark EMP (1981) *Principles and Applications of Medicolegal Alcohol Determination*. Davis (ed.). Biomedical Publications.
- Womelsdorf T, Vinck M, Leung LS, et al. (2010) Selective theta-synchronization of choice-relevant information subserves goal-directed behavior. *Frontiers in human neuroscience* 4: 210. DOI: 10.3389/fnhum.2010.00210.
- Zuckerman M and Link K (1968) Construct validity for the sensation-seeking scale. *Journal of*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

consulting and clinical psychology 32(4): 420–6.




Figure 1. Schematic diagram of the task. The stimuli were presented for 100 ms every 1400±200 ms.




Figure 2. Group-average difference maps (LD minus BD) of event-related theta source power. Time courses show theta power estimated to the anterior (aIFC) and posterior (pIFC) inferior frontal cortices, the sensorimotor (sMOT) areas and the occipital cortex (Occ) expressed as percent change from baseline. Targets evoked greater theta activity overall which was especially prominent in the rIFC. Theta power in the rIFC was reduced in BD compared to LD participants. Light-shaded vertical bars indicate the analyzed time windows: 50-150 ms (Occ) and the 380-510 ms for the anterior ROIs.




Figure 3. Scatter plots of the correlations between target theta power in the right aIFC and the performance accuracy, D2 attentional test scores, the estimated BAC and the AUDIT scores.

PLV - Theta band (4-7 Hz) at 410-580 ms





Figure 4. Group average maps of event-related theta source estimates and timecourses of phase-locking values (PLV) calculated between the aIFC and the pIFC. The PLVs are expressed as percent change from baseline. The anterior (aIFC) and posterior areas (pIFC) of the right IFC showed increased oscillatory synchrony in LD from 410-580 ms where PLV decreased in BD and correlated negatively with alcohol consumption (est-BAC).



33 additional figure for reviewer 1. PLV test